

Electromagnetic fields act *via* activation of voltage-gated calcium channels to produce beneficial or adverse effects

1. Martin L. Pall*

Article first published online: 26 JUN 2013

DOI: 10.1111/jcmm.12088

© 2013 The Author. Journal of Cellular and Molecular Medicine Published by Foundation for Cellular and Molecular Medicine/Blackwell Publishing Ltd

This is an open access article under the terms of the [Creative Commons Attribution](#) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Available on the publisher's web site:

<http://onlinelibrary.wiley.com/doi/10.1111/jcmm.12088/pdf>



Journal of Cellular and Molecular Medicine

Keywords:

- intracellular Ca²⁺;
- voltage-gated calcium channels;
- low frequency electromagnetic field exposure;
- nitric oxide;
- oxidative stress;
- calcium channel blockers

Abstract

The direct targets of extremely low and microwave frequency range electromagnetic fields (EMFs) in producing non-thermal effects have not been clearly established. However, studies in the literature, reviewed here, provide substantial support for such direct targets. Twenty-three studies have shown that voltage-gated calcium channels (VGCCs) produce these and other EMF effects, such that the L-type or other VGCC blockers block or greatly lower diverse EMF effects. Furthermore, the voltage-gated properties of these channels may provide biophysically plausible mechanisms for EMF biological effects. Downstream responses of such EMF exposures may be mediated through Ca²⁺/calmodulin stimulation of nitric oxide synthesis. Potentially, physiological/therapeutic responses may be largely as a result of nitric oxide-cGMP-protein kinase G pathway stimulation. A well-studied example of such an apparent therapeutic response, EMF stimulation of bone growth, appears to work along this pathway. However, pathophysiological responses to EMFs may be as a result of nitric oxide-peroxynitrite-oxidative stress pathway of action. A single such well-documented example, EMF induction of DNA single-strand breaks in cells, as measured by alkaline comet assays, is reviewed here. Such single-strand breaks are known to be produced through the action of this pathway. Data on the mechanism of EMF induction of such breaks are limited; what data are available support this proposed mechanism. Other Ca²⁺-mediated regulatory changes, independent of nitric oxide, may also have roles. This article reviews, then, a substantially supported set of targets, VGCCs, whose stimulation produces non-thermal EMF responses by humans/higher animals with

downstream effects involving Ca²⁺/calmodulin-dependent nitric oxide increases, which may explain therapeutic and pathophysiological effects.

Introduction

An understanding of the complex biology of the effects of electromagnetic fields (EMFs) on human/higher animal biology inevitably must be derived from an understanding of the target or targets of such fields in the impacted cells and tissues. Despite this, no understanding has been forthcoming on what those targets are and how they may lead to the complex biological responses to EMFs composed of low-energy photons. The great puzzle, here, is that these EMFs are comprised of low-energy photons, those with insufficient energy to individually influence the chemistry of the cell, raising the question of how non-thermal effects of such EMFs can possibly occur. The author has found that there is a substantial literature possibly pointing to the direct targets of such EMFs and it is the goal of this study to review that evidence as well as review how those targets may lead to the complex biology of EMF exposure.

The role of increased intracellular Ca²⁺ following EMF exposure was already well documented more than 20 years ago, when Walleczek [1] reviewed the role of changes in calcium signalling that were produced in response EMF exposures. Other, more recent studies have confirmed the role of increased intracellular Ca²⁺ following EMF exposure, a few of which are discussed below. His review [1] included two studies [2, 3] that showed that the L-type voltage-gated channel blocker, verapamil could lower or block changes in response to EMFs. The properties of voltage-gated calcium channels (VGCCs) have been reviewed elsewhere [4]. Subsequently, extensive evidence has been published clearly showing that the EMF exposure can act to produce excessive activity of the VGCCs in many cell types [5-26] suggesting that these may be direct targets of EMF exposure. Many of these studies implicate specifically the L-type VGCCs such that various L-type calcium channel blockers can block responses to EMF exposure (Table 1). However, other studies have shown lowered responses produced by other types of calcium channel blockers including N-type, P/Q-type, and T-type blockers (Table 1), showing that other VGCCs may have important roles. Diverse responses to EMFs are reported to be blocked by such calcium channel blockers (Table 1), suggesting that most if not all EMF-mediated responses may be produced through VGCC stimulation. Voltage-gated calcium channels are essential to the responses produced by extremely low frequency (including 50/60 Hz) EMFs and also to microwave frequency range EMFs, nanosecond EMF pulses, and static electrical and magnetic fields (Table 1).

Table 1. EMF responses blocked or lowered by calcium channel blockers

Ref. no.	EMF type	Calcium channel	Cell type or organism	Response measured
1. EMF: electromagnetic field; ELF: extremely low frequency.				
[2]	Pulsed magnetic fields	L-type	Human lymphocytes	Cell proliferation; cytokine production
[3]	Static magnetic field (0.1 T)	L-type	Human polymorphonuclear leucocytes	Cell migration; degranulation
[5]	ELF	L-type	Rat chromaffin cells	Differentiation; catecholamine release
[6]	Electric field	L-type	Rat and mouse bone cells	Increased Ca ²⁺ , phospholipase A2, PGE2
[7]	50 Hz	L-type	Mytilus (mussel) immunocytes	Reduced shape change, cytotoxicity
[8]	50 Hz	L-type	AtT20 D16V, mouse pituitary corticotrope-derived	Ca ²⁺ increase; cell morphology, premature differentiation
[9]	50 Hz	L-type	Neural stem/progenitor cells	<i>In vitro</i> differentiation, neurogenesis
[10]	Static magnetic field	L-type	Rat	Reduction in oedema formation

Table 1. EMF responses blocked or lowered by calcium channel blockers

Ref. no.	EMF type	Calcium channel	Cell type or organism	Response measured
[11]	NMR	L-type	Tumour cells	Synergistic effect of EMF on anti-tumour drug toxicity
[12]	Static magnetic field	L-type	Myelomonocytic U937 cells	Ca ²⁺ influx into cells and anti-apoptotic effects
[13]	60 Hz	L-type	Mouse	Hyperalgesic response to exposure
[14]	Single nanosecond electric pulse	L-type	Bovine chromaffin cells	Very rapid increase in intracellular Ca ²⁺
[15]	Biphasic electric current	L-type	Human mesenchymal stromal cells	Osteoblast differentiation and cytokine production
[16]	DC & AC magnetic fields	L-type	β-cells of pancreas, patch clamped	Ca ²⁺ flux into cells
[17]	50 Hz	L-type	Rat pituitary cells	Ca ²⁺ flux into cells
[18]	50 Hz	L-type, N-type	Human neuroblastoma IMR32 and rat pituitary GH3 cells	Anti-apoptotic activity
[19]	Nanosecond pulse	L-type, N-type, P/Q-type	Bovine chromaffin cells	Ca ²⁺ dynamics of cells
[20]	50 Hz	Not determined	Rat dorsal root ganglion cells	Firing frequency of cells
[21]	700–1100 MHz	N-type	Stem cell–derived neuronal cells	Ca ²⁺ dynamics of cells
[22]	Very weak electrical fields	T-type	Sharks	Detection of very weak magnetic fields in the ocean
[23]	Short electric pulses	L-type	Human eye	Effect on electro-oculogram
[24]	Weak static magnetic field	L-type	Rabbit	Baroreflex sensitivity
[25]	Weak electric fields	T-type	Neutrophils	Electrical and ion dynamics
[26]	Static electric fields, ‘capacitive’	L-type	Bovine articular chondrocytes	Aggrecan & type II collagen expression; calcineurin and other Ca ²⁺ /calmodulin responses

In a recent study, Pilla [27] showed that an increase in intracellular Ca²⁺ must have occurred almost immediately after EMF exposure, producing a Ca²⁺/calmodulin-dependent increase in nitric oxide occurring in less than 5 sec. Although Pilla [27] did not test whether VGCC stimulation was involved in his study, there are few alternatives that can produce such a rapid Ca²⁺ response, none of which has been implicated in EMF responses. Other studies, each involving VGCCs, summarized in Table 1, also showed rapid Ca²⁺ increases following EMF exposure [8, 16, 17, 19, 21]. The rapidity of these responses rule out many types of regulatory interactions as being involved in producing the increased VGCC activity following EMF exposure and suggests, therefore, that VGCC stimulation in the plasma membrane is directly produced by EMF exposure.

Possible modes of action following VGCC stimulation

The increased intracellular Ca²⁺ produced by such VGCC activation may lead to multiple regulatory responses, including the increased nitric oxide levels produced through the action of the two

Ca²⁺/calmodulin-dependent nitric oxide synthases, nNOS and eNOS. Increased nitric oxide levels typically act in a physiological context through increased synthesis of cGMP and subsequent activation of protein kinase G [28, 29]. In contrast, in most pathophysiological contexts, nitric oxide reacts with superoxide to form peroxynitrite, a potent non-radical oxidant [30, 31], which can produce radical products, including hydroxyl radical and NO₂ radical [32].

Therapeutic bone-growth stimulation *via* Ca²⁺/nitric oxide/cGMP/protein kinase G

An example of a therapeutic effect for bone repair of EMF exposure in various medical situations includes increasing osteoblast differentiation and maturation and has been reviewed repeatedly [33-44]. The effects of EMF exposure on bone cannot be challenged, although there is still considerable question about the best ways to apply this clinically [33-44]. Our focus, here, is to consider possible mechanisms of action.

Multiple studies have implicated increased Ca²⁺ and nitric oxide in the EMF stimulation of bone growth [44-49]; three have also implicated increased cGMP and protein kinase G activity [46, 48, 49]. In addition, studies on other regulatory stimuli leading to increased bone growth have also implicated increased cGMP levels and protein kinase G in this response [50-56]. In summary, then, it can be seen from the above that there is a very well-documented action of EMFs in stimulating osteoblasts and bone growth. The available data, although limited, support the action of the main pathway involved in physiological responses to Ca²⁺ and nitric oxide, namely Ca²⁺/nitric oxide/cGMP/protein kinase G in producing such stimulation.

Ca²⁺/nitric oxide/peroxynitrite and pathophysiological responses to EMF exposures: the example of single-strand DNA breaks

As was noted above, most of the pathophysiological effects of nitric oxide are mediated through peroxynitrite elevation and consequent oxidative stress. There are many reviews and other studies, implicating oxidative stress in generating pathophysiological effects of EMF exposure [see for example 57-64]. In some of these studies, the rise in oxidative stress markers parallels the rise in nitric oxide, suggesting a peroxynitrite-mediated mechanism [64-67].

Peroxyntirite elevation is usually measured through a marker of peroxynitrite-mediated protein nitration, 3-nitrotyrosine (3-NT). There are four studies where 3-NT levels were measured before and after EMF exposure [66, 68-70]. Each of these studies provides some evidence supporting the view that EMF exposure increases levels of peroxynitrite and therefore 3-NT levels [66, 68-70]. Although these cannot be taken as definitive, when considered along with the evidence on oxidative stress and elevated nitric oxide production in response to EMF exposure, they strongly suggest a peroxynitrite-mediated mechanism of oxidative stress in response to EMFs.

Such a peroxynitrite-mediated mechanism may explain the many studies showing the single-stranded breaks in DNA, as shown by alkaline comet assays or the similar microgel electrophoresis assay, following EMF exposures in most such studies [71-89], but not in all [90-97]. Some of the factors that are reported to influence whether such DNA single-strand breaks are detected after EMF exposure include the type of cell studied [79, 86], dosage of EMF exposure [78] and the type of EMF exposure studied [73, 77]. Oxidative stress and free radicals have roles, both because there is a concomitant increase in oxidative stress and because antioxidants have been shown to greatly lower the generation of DNA single-strand breaks following EMF exposure [72, 75, 81, 82] as has also been shown for peroxynitrite-mediated DNA breaks produced under other conditions. It has also been shown that one can block the generation of DNA single-strand breaks with a nitric oxide synthase inhibitors [82].

Peroxyntirite has been shown to produce single-strand DNA breaks [98-100], a process that is inhibited by many but not all antioxidants [99, 100]. It can be seen from this that the data on generation of single-strand DNA breaks, although quite limited, support a mechanism involving nitric oxide/peroxynitrite/free radical (oxidative stress). Although the data on the possible role of peroxynitrite in EMF-induced DNA single-strand breaks are limited, what data are available supports such a peroxynitrite role.

Discussion and conclusions

How do EMFs composed of low-energy photons produce non-thermal biological changes, both pathophysiological and, in some cases, potentially therapeutic, in humans and higher animals? It may be surprising that the answer to this question has been hiding in plain sight in the scientific literature.

However, in this era of highly focused and highly specialized science, few of us have the time to read the relevant literature, let alone organize the information found within it in useful and critical ways.

This study shows that:

1. Twenty-three different studies have found that such EMF exposures act *via* activation of VGCCs, such that VGCC channel blockers can prevent responses to such exposures (Table 1). Most of the studies implicate L-type VGCCs in these responses, but there are also other studies implicating three other classes of VGCCs.
2. Both extremely low frequency fields, including 50/60 cycle exposures, and microwave EMF range exposures act *via* activation of VGCCs. So do static electric fields, static magnetic fields and nanosecond pulses.
3. Voltage-gated calcium channel stimulation leads to increased intracellular Ca^{2+} , which can act in turn to stimulate the two calcium/calmodulin-dependent nitric oxide synthases and increase nitric oxide. It is suggested here that nitric oxide may act in therapeutic/potentially therapeutic EMF responses *via* its main physiological pathway, stimulating cGMP and protein kinase G. It is also suggested that nitric oxide may act in pathophysiological responses to EMF exposure, by acting as a precursor of peroxynitrite, producing both oxidative stress and free radical breakdown products.
4. The interpretation in three above is supported by two specific well-documented examples of EMF effects. Electromagnetic fields stimulation of bone growth, modulated through EMF stimulation of osteoblasts, appears to involve an elevation/nitric oxide/protein kinase G pathway. In contrast to that, it seems likely that the EMF induction of single-stranded DNA breaks involves a Ca^{2+} /elevation/nitric oxide/peroxynitrite/free radical (oxidative stress) pathway.

It may be asked why we have evidence for involvement of VGCCs in response to EMF exposure, but no similar evidence for involvement of voltage-gated sodium channels? Perhaps, the reason is that there are many important biological effects produced in increased intracellular Ca^{2+} , including but not limited to nitric oxide elevation, but much fewer are produced by elevated Na^+ .

The possible role of peroxynitrite as opposed to protein kinase G in producing pathophysiological responses to EMF exposure raises the question of whether there are practical approaches to avoiding such responses? Typically peroxynitrite levels can be highly elevated when both of its precursors, nitric oxide and superoxide, are high. Consequently, agents that lower nitric oxide synthase activity and agents that raise superoxide dismutases (SODs, the enzymes that degrade superoxide) such as phenolics and other Nrf2 activators that induce SOD activity [101], as well as calcium channel blockers may be useful. Having said that, this is a complex area, where other approaches should be considered, as well.

Although the various EMF exposures as well as static electrical field exposures can act to change the electrical voltage-gradient across the plasma membrane and may, therefore, be expected to stimulate VGCCs through their voltage-gated properties, it may be surprising that static magnetic fields also act to activate VGCCs because static magnetic fields do not induce electrical changes on static objects. However, cells are far from static. Such phenomena as cell ruffling [101,101] may be relevant, where thin cytoplasmic sheets bounded on both sides by plasma membrane move rapidly. Such rapid movement of the electrically conducting cytoplasm, may be expected to influence the electrical charge across the plasma membrane, thus potentially stimulating the VGCCs.

Earlier modelling of electrical effects across plasma membranes of EMF exposures suggested that such electrical effects were likely to be too small to explain EMF effects at levels reported to produce biological changes (see, for example [22]). However, more recent and presumably more biologically plausible modelling have suggested that such electrical effects may be much more substantial [104-109] and may, therefore, act to directly stimulate VGCCs.

Direct stimulation of VGCCs by partial depolarization across the plasma membrane is suggested by the following observations discussed in this review:

1. The very rapid, almost instantaneous increase in intracellular Ca^{2+} found in some studies following EMF exposure [8, 16, 17, 19, 21, 27]. The rapidity here means that most, if not all indirect, regulatory effects can be ruled out.
2. The fact that not just L-type, but three additional classes of VGCCs are implicated in generating biological responses to EMF exposure (Table 1), suggesting that their voltage-gated properties may be a key feature in their ability to respond to EMFs.
3. Most, if not all, EMF effects are blocked by VGCC channel blockers (Table 1).
4. Modelling of EMF effects on living cells suggests that plasma membrane voltage changes may have key roles in such effects [104-109]. Saunders and Jefferys stated [110] that 'It is well established that electric fields ... or exposure to low frequency magnetic fields, will, if of sufficient magnitude, excite nerve tissue through their interactions with ... voltage gated ion

channels'. They further state [110] that this is achieved by direct effects on the electric dipole voltage sensor within the ion channel.

One question that is not answered by any of the available data is whether what is known as 'dirty electricity' [111-113], generated by rapid, in many cases, square wave transients in EMF exposure, also acts by stimulating VGCCs. Such dirty electricity is inherent in any digital technology because digital technology is based on the use of such square wave transients and it may, therefore, be of special concern in this digital era, but there have been no tests of such dirty electricity that determine whether VGCCs have roles in response to such fields, to my knowledge. The nanosecond pulses, which are essentially very brief, but high-intensity dirty electricity do act, at least in part, *via* VGCC stimulation (Table 1), suggesting that dirty electricity may do likewise. Clearly, we need direct study of this question.

The only detailed alternative to the mechanism of non-thermal EMF effects discussed here, to my knowledge, is the hypothesis of Friedman *et al.* [114] and supported by Desai *et al.* [115] where the apparent initial response to EMF exposure was proposed to be NADH oxidase activation, leading to oxidative stress and downstream regulatory effects. Although they provide some correlative evidence for a possible role of NADH oxidase [114], the only causal evidence is based on a presumed specific inhibitor of NADH oxidase, diphenyleneiodonium (DPI). However, DPI has been shown to be a non-specific cation channel blocker [116], clearly showing a lack of such specificity and suggesting that it may act, in part, as a VGCC blocker. Consequently, a causal role for NADH oxidase in responses to EMF exposure must be considered to be undocumented.

In summary, the non-thermal actions of EMFs composed of low-energy photons have been a great puzzle, because such photons are insufficiently energetic to directly influence the chemistry of cells. The current review provides support for a pathway of the biological action of ultralow frequency and microwave EMFs, nanosecond pulses and static electrical or magnetic fields: EMF activation of VGCCs leads to rapid elevation of intracellular Ca^{2+} , nitric oxide and in some cases at least, peroxynitrite. Potentially therapeutic effects may be mediated through the Ca^{2+} /nitric oxide/cGMP/protein kinase G pathway. Pathophysiological effects may be mediated through the Ca^{2+} /nitric oxide/peroxynitrite pathway. Other Ca^{2+} -mediated effects may have roles as well, as suggested by Xu *et al.* [26].

Conflicts of interest

The author confirms that there are no conflicts of interest.

References

- 1
Walleczek J. *Electromagnetic field effects on cells of the immune system: the role of calcium signaling*. *FASEB J.* 1992; 6: 3177–85.
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 213](#)
- 2
Cadossi R, Emilia G, Ceccherelli G, *et al.* 1988 *Lymphocytes and pulsing magnetic fields*. In: Marino EE, editor. *Modern bioelectricity*. New York: Dekker; 1998. pp. 451–96.
- 3
Papatheofanis FJ. *Use of calcium channel antagonists as magnetoprotective agents*. *Radiat Res.* 1990; 122: 24–8.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 19](#)
- 4
Catterall WA. *Structure and regulation of voltage-gated Ca^{2+} channels*. *Annu Rev Cell Dev Biol.* 2000; 16: 521–55.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 1224](#)
- 5

- Morgado-Valle C, Verdugo-Díaz L, García DE, et al. The role of voltage-gated Ca²⁺ channels in neurite growth of cultured chromaffin cells induced by extremely low frequency (ELF) magnetic field stimulation. Cell Tissue Res. 1998; 291: 217–30.*
- [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 40](#)
- 6
- Lorich DG, Brighton CT, Gupta R, et al. Biochemical pathway mediating the response of bone cells to capacitive coupling. Clin Orthop Relat Res. 1998; 246–56.*
- [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 30](#)
- 7
- Gobba F, Malagoli D, Ottaviani E. Effects of 50 Hz magnetic fields on fMLP-induced shape changes in invertebrate immunocytes: the role of calcium ion channels. Bioelectromagnetics. 2003; 24: 277–82.*
- Direct Link:
- [Abstract](#)
 - [PDF\(119K\)](#)
 - [References](#)
 - [Web of Science® Times Cited: 9](#)
- 8
- Lisi A, Ledda M, Rosola E, et al. Extremely low frequency electromagnetic field exposure promotes differentiation of pituitary corticotrope-derived AtT20 D16V cells. Bioelectromagnetics. 2006; 27: 641–51.*
- Direct Link:
- [Abstract](#)
 - [PDF\(433K\)](#)
 - [References](#)
 - [Web of Science® Times Cited: 14](#)
- 9
- Piacentini R, Ripoli C, Mezzogori D, et al. Extremely low-frequency electromagnetic fields promote in vitro neurogenesis via upregulation of Ca(v)1-channel activity. J Cell Physiol. 2008; 215: 129–39.*
- Direct Link:
- [Abstract](#)
 - [Full Article \(HTML\)](#)
 - [PDF\(415K\)](#)
 - [References](#)
 - [Web of Science® Times Cited: 36](#)
- 10
- Morris CE, Skalak TC. Acute exposure to a moderate strength static magnetic field reduces edema formation in rats. Am J Physiol Heart Circ Physiol. 2008; 294: H50–7.*
- [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 14](#)
- 11
- Ghibelli L, Cerella C, Cordisco S, et al. NMR exposure sensitizes tumor cells to apoptosis. Apoptosis. 2006; 11: 359–65.*
- [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 10](#)

- 12
Fanelli C, Coppola S, Barone R, et al. Magnetic fields increase cell survival by inhibiting apoptosis via modulation of Ca²⁺ influx. FASEB J. 1999; 13: 95–102.
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 115](#)
- 13
Jeong JH, Kum C, Choi HJ, et al. Extremely low frequency magnetic field induces hyperalgesia in mice modulated by nitric oxide synthesis. Life Sci. 2006; 78: 1407–12.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 16](#)
- 14
Vernier PT, Sun Y, Chen MT, et al. Nanosecond electric pulse-induced calcium entry into chromaffin cells. Bioelectrochemistry. 2008; 73: 1–4.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 17](#)
- 15
Kim IS, Song JK, Song YM, et al. Novel effect of biphasic electric current on in vitro osteogenesis and cytokine production in human mesenchymal stromal cells. Tissue Eng Part A. 2009; 15: 2411–22.
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 18](#)
- 16
Höjveik P, Sandblom J, Galt S, et al. Ca²⁺ ion transport through patch-clamped cells exposed to magnetic fields. Bioelectromagnetics. 1995; 16: 33–40.
 Direct Link:
 - [Abstract](#)
 - [PDF\(683K\)](#)
 - [References](#)
 - [Web of Science® Times Cited: 27](#)
- 17
Barbier E, Vetret B, Dufy B. Stimulation of Ca²⁺ influx in rat pituitary cells under exposure to a 50 Hz magnetic field. Bioelectromagnetics. 1996; 17: 303–11.
 Direct Link:
 - [Abstract](#)
 - [PDF\(881K\)](#)
 - [References](#)
 - [Web of Science® Times Cited: 41](#)
- 18
Grassi C, D'Ascenzo M, Torsello A, et al. Effects of 50 Hz electromagnetic fields on voltage-gated Ca²⁺ channels and their role in modulation of neuroendocrine cell proliferation and death. Cell Calcium. 2004; 35: 307–15.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 40](#)
- 19
Craviso GL, Choe S, Chatterjee P, et al. Nanosecond electric pulses: a novel stimulus for triggering Ca²⁺ influx into chromaffin cells via voltage-gated Ca²⁺ channels. Cell Mol Neurobiol. 2010; 30: 1259–65.

- [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 8](#)
- 20

Marchionni I, Paffi A, Pellegrino M, et al. Comparison between low-level 50 Hz and 900 MHz electromagnetic stimulation on single channel ionic currents and on firing frequency in dorsal root ganglion isolated neurons. Biochim Biophys Acta. 2006; 1758: 597–605.

 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 11](#)
- 21

Rao VS, Titushkin IA, Moros EG, et al. Nonthermal effects of radiofrequency-field exposure on calcium dynamics in stem cell-derived neuronal cells: elucidation of calcium pathways. Radiat Res. 2008; 169: 319–29.

 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 10](#)
- 22

Adair RK, Astumian RD, Weaver JC. Detection of weak electric fields by sharks, rays and skates. Chaos. 1998; 8: 576–87.

 - [CrossRef](#),
 - [Web of Science® Times Cited: 25](#),
 - [ADS](#)
- 23

Constable PA. Nifedipine alters the light-rise of the electro-oculogram in man. Graefes Arch Clin Exp Ophthalmol. 2011; 249: 677–84.

 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 1](#)
- 24

Gmitrov J, Ohkuba C. Verapamil protective effect on natural and artificial magnetic field cardiovascular impact. Bioelectromagnetics. 2002; 23: 531–41.

Direct Link:

 - [Abstract](#)
 - [PDF\(171K\)](#)
 - [References](#)
 - [Web of Science® Times Cited: 19](#)
- 25

Kindzelskii AL, Petty HR. Ion channel clustering enhances weak electric field detection by neutrophils: apparent role of SKF96365-sensitive cation channels and myeloperoxidase trafficking cellular responses. Eur Biophys J. 2005; 35: 1–26.

 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 17](#)
- 26

Xu J, Wang W, Clark CC, et al. Signal transduction in electrically stimulated articular chondrocytes involves translocation of extracellular calcium through voltage-gated channels. Osteoarthritis Cartilage. 2009; 17: 397–405.

 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 6](#)

- 27
Pilla AA. *Electromagnetic fields instantaneously modulate nitric oxide signaling in challenged biological systems. Biochem Biophys Res Commun.* 2012; 426: 330–3.
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 3](#)
- 28
McDonald LJ, Murad F. *Nitric oxide and cyclic GMP signaling. Proc Soc Exp Biol Med.* 1996; 211: 1–6.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#)
- 29
Francis SH, Busch JL, Corbin JD, et al. *cGMP-dependent protein kinases and cGMP phosphodiesterases in nitric oxide and cGMP action. Pharmacol Rev.* 2010; 62: 525–63.
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 93](#)
- 30
Pacher P, Beckman JS, Liaudet L. *Nitric oxide and peroxynitrite in health and disease. Physiol Rev.* 2007; 87: 315–424.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 1189](#)
- 31
Pryor WA, Squadrito GL. *The chemistry of peroxynitrite: a product from the reaction of nitric oxide with superoxide. Am J Physiol.* 1995; 268: L699–722.
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 889](#)
- 32
Lymar SV, Khairutdinov RF, Hurst JK. *Hydroxyl radical formation by O-O bond homolysis in peroxynitrous acid. Inorg Chem.* 2003; 42: 5259–66.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 48](#)
- 33
Ryabi JT. *Clinical effects of electromagnetic fields on fracture healing. Clin Orthop Relat Res.* 1998; 355(Suppl. 1): S205–15.
 - [CrossRef](#)
- 34
Oishi M, Onesti ST. *Electrical bone graft stimulation for spinal fusion: a review. Neurosurgery.* 2000; 47: 1041–55.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 11](#)
- 35
Aaron RK, Ciombor DM, Simon BJ. *Treatment of nonunions with electric and electromagnetic fields. Clin Orthop Relat Res.* 2004; 10: 579–93.
- 36
Goldstein C, Sprague S, Petrisor BA. *Electrical stimulation for fracture healing: current evidence. J Orthop Trauma.* 2010; 24(Suppl. 1): S62–5.

- [CrossRef](#),
 - [Web of Science® Times Cited: 10](#)
- 37
Demetriou R, Babis GC. Biomaterial osseointegration enhancement with biophysical stimulation. J Musculoskelet Neuronal Interact. 2007; 7: 253–65.
- 38
Griffin XL, Warner F, Costa M. The role of electromagnetic stimulation in the management of established non-union of long bone fractures: what is the evidence? Injury. 2008; 39: 419–29.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 25](#)
- 39
Huang LQ, He HC, He CQ, et al. Clinical update of pulsed electromagnetic fields on osteoporosis. Chin Med J. 2008; 121: 2095–9.
 - [PubMed](#),
 - [Web of Science® Times Cited: 4](#)
- 40
Groah SL, Lichy AM, Libin AV, et al. Intensive electrical stimulation attenuates femoral bone loss in acute spinal cord injury. PM R. 2010; 2: 1080–7.
 - [CrossRef](#),
 - [Web of Science® Times Cited: 4](#)
- 41
Schidt-Rohlfing B, Silny J, Gavenis K, et al. Electromagnetic fields, electric current and bone healing – what is the evidence? Z Orthop Unfall. 2011; 149: 265–70.
- 42
Griffin XL, Costa ML, Parsons N, et al. Electromagnetic field stimulation for treating delayed union or non-union of long bone fractures in adults. Cochrane Database Syst Rev. 2011; CDO08471. doi: 10.1002/14651858.CD008471.pub2.
- 43
Chalidis B, Sachinis N, Assiotis A, et al. Stimulation of bone formation and fracture healing with pulsed electromagnetic fields: biologic responses and clinical implications. Int J Immunopathol Pharmacol. 2011; 24(1 Suppl. 2): 17020.
- 44
Zhong C, Zhao TF, Xu ZJ, et al. Effects of electromagnetic fields on bone regeneration in experimental and clinical studies: a review of the literature. Chin Med J. 2012; 125: 367–72.
 - [CAS](#),
 - [Web of Science®](#)
- 45
Diniz P, Soejima K, Ito G. Nitric oxide mediates the effects of pulsed electromagnetic field stimulation on the osteoblast proliferation and differentiation. Nitric Oxide. 2002; 7: 18–23.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 36](#)
- 46
Fitzsimmons RJ, Gordon SL, Ganey T, et al. A pulsing electric field (PEF) increases human chondrocyte proliferation through a transduction pathway involving nitric oxide signaling. J Orthopaedic Res. 2008; 26: 854–9.
Direct Link:
 - [Abstract](#)
 - [PDF\(174K\)](#)
 - [References](#)
 - [Web of Science® Times Cited: 17](#)
- 47

- Lin H-Y, Lin Y-J. *In vitro effects of low frequency electromagnetic fields on osteoblast proliferation and maturation in an inflammatory environment. Bioelectromagnetics. 2011; 32: 552–60.*
- Direct Link:
- [Abstract](#)
 - [Full Article \(HTML\)](#)
 - [PDF\(2955K\)](#)
 - [References](#)
 - [Web of Science® Times Cited: 6](#)
- 48
Cheng G, Zhai Y, Chen K, et al. *Sinusoidal electromagnetic field stimulates rat osteoblast differentiation and maturation via activation of NO-cGMP-PKG pathway. Nitric Oxide. 2011; 25: 316–25.*
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 3](#)
 - 49
Pilla A, Fitzsimmons R, Muehsam D, et al. *Electromagnetic fields as first messenger in biological signaling: application to calmodulin-dependent signaling in tissue repair. Biochim Biophys Acta. 2011; 1810: 1236–45.*
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 7](#)
 - 50
Rangaswami H, Schwappacher R, Tran T, et al. *Protein kinase G and focal adhesion kinase converge on Src/Akt/ β -catenin signaling module in osteoblast mechanotransduction. J Biol Chem. 2012; 287: 21509–19.*
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 1](#)
 - 51
Marathe N, Rangaswami H, Zhuang S, et al. *Pro-survival effects of 17 β -estradiol on osteocytes are mediated by nitric oxide/cGMP via differential actions of cGMP-dependent protein kinases I and II. J Biol Chem. 2012; 287: 978–88.*
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 3](#)
 - 52
Rangaswami H, Schwappacher R, Marathe N, et al. *Cyclic GMP and protein kinase G control a Src-containing mechanosome in osteoblasts. Sci Signal. 2010; 3: ra91.*
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 17](#)
 - 53
Rangaswami H, Marathe N, Zhuang S, et al. *Type II cGMP-dependent protein kinase mediates osteoblast mechanotransduction. J Biol Chem. 2009; 284: 14796–808.*
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 22](#)
 - 54
Saura M, Tarin C, Zaragoza C. *Recent insights into the implication of nitric oxide in osteoblast differentiation and proliferation during bone development. ScientificWorldJournal. 2010; 10: 624–32.*
 - [CrossRef](#),

- [CAS](#),
 - [Web of Science® Times Cited: 10](#)
- 55
Zaragoza C, López-Rivera E, García-Rama C, et al. Cbfa-1 mediates nitric oxide regulation of MMP-13 in osteoblasts. J Cell Sci. 2006; 119: 1896–902.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 21](#)
- 56
Wang DH, Hu YS, Du JJ, et al. Ghrelin stimulates proliferation of human osteoblastic TE85 cells via NO/cGMP signaling pathway. Endocrine. 2009; 35: 112–7.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 12](#)
- 57
Simkó M. Cell type specific redox status is responsible for diverse electromagnetic field effects. Curr Med Chem. 2007; 14: 1141–52.
 - [CrossRef](#),
 - [PubMed](#),
 - [Web of Science® Times Cited: 46](#)
- 58
Consales C, Merla C, Marino C, et al. Electromagnetic fields, oxidative stress, and neurodegeneration. Int J Cell Biol. 2012; 2012: 683897.
- 59
Johansson O. Disturbance of the immune system by electromagnetic fields—A potentially underlying cause for cellular damage and tissue repair reduction which could lead to disease and impairment. Pathophysiology. 2009; 16: 157–77.
 - [CrossRef](#),
 - [PubMed](#)
- 60
Kovacic P, Somanathan R. Electromagnetic fields: mechanism, cell signaling, other bioprocesses, toxicity, radicals, antioxidants and beneficial effects. J Recept Signal Transduct Res. 2010; 30: 214–26.
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 7](#)
- 61
Wolf FI, Torsello A, Tedesco B, et al. 50-Hz extremely low frequency electromagnetic fields enhance cell proliferation and DNA damage: possible involvement of a redox mechanism. Biochim Biophys Acta. 2005; 1743: 120–9.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 98](#)
- 62
Iakimenko IL, Sidorik EP, Tsybulin AS. Metabolic changes in cells under electromagnetic radiation of mobile communication systems. Ukr Biokhim Zh. 2011; 83: 20–8.
 - [CAS](#)
- 63
Jing J, Yuhua Z, Xiao-qian Y, et al. The influence of microwave radiation from cellular phone on fetal rat brain. Electromagn Biol Med. 2012; 31: 57–66.
 - [CrossRef](#),
 - [CAS](#),

- [Web of Science®](#)
- 64
Esmekaya MA, Ozer C, Seyhan N. 900 MHz pulse-modulated radiofrequency radiation induces oxidative stress on heart, lung, testis and liver tissues. Gen Physiol Biophys. 2011; 30: 84–9.
 - [CrossRef](#),
 - [Web of Science® Times Cited: 3](#)
- 65
Aydin B, Akar A. Effects of a 900-MHz electromagnetic field on oxidative stress parameters in rat lymphoid organs, polymorphonuclear leukocytes and plasma. Arch Med Res. 2011; 42: 261–7.
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 6](#)
- 66
Guler G, Turkozer Z, Tomruk A, et al. The protective effects of N-acetyl-L-cysteine and epigallocatechin-3-gallate on electric field-induced hepatic oxidative stress. Int J Radiat Biol. 2008; 84: 669–80.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 7](#)
- 67
Guney M, Ozguner F, Oral B, et al. 900 MHz radiofrequency-induced histopathologic changes and oxidative stress in rat endometrium: protection by vitamins E and C. Toxicol Ind Health. 2007; 23: 411–20.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 24](#)
- 68
Sypniewska RK, Millenbaugh NJ, Kiel JL, et al. Protein changes in macrophages induced by plasma from rats exposed to 35 GHz millimeter waves. Bioelectromagnetics. 2010; 31: 656–63.
 Direct Link:
 - [Abstract](#)
 - [Full Article \(HTML\)](#)
 - [PDF\(201K\)](#)
 - [References](#)
 - [Web of Science® Times Cited: 1](#)
- 69
Grigoriev YG, Mikhailov VF, Ivanov AA, et al. Autoimmune processes after long-term low-level exposure to electromagnetic fields part 4. Oxidative intracellular stress response to the long-term rat exposure to nonthermal RF EMF. Biophysics. 2010; 55: 1054–8.
 - [CrossRef](#)
- 70
Erdal N, Gürgül S, Tamer L, et al. Effects of long-term exposure of extremely low frequency magnetic field on oxidative/nitrosative stress in rat liver. J Radiat Res. 2008; 49: 181–7.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 9](#)
- 71
Ahuja YR, Vijayashree B, Saran R, et al. In vitro effects of low-level, low-frequency electromagnetic fields on DNA damage in human leucocytes by comet assay. Indian J Biochem Biophys. 1999; 36: 318–22.
 - [PubMed](#),
 - [CAS](#),

- [Web of Science® Times Cited: 14](#)
- 72
Amara S, Douki T, Ravanat JL, et al. Influence of a static magnetic field (250 mT) on the antioxidant response and DNA integrity in THP1 cells. Phys Med Biol. 2007; 52: 889–98.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 21](#),
 - [ADS](#)
- 73
Focke F, Schuermann D, Kuster N, et al. DNA fragmentation in human fibroblasts under extremely low frequency electromagnetic field exposure. Mutat Res. 2010; 683: 74–83.
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 24](#)
- 74
Franzellitti S, Valbonesi P, Ciancaglini N, et al. Transient DNA damage induced by high-frequency electromagnetic fields (GSM 1.8 GHz) in the human trophoblast HTR-8/SVneo cell line evaluated with the alkaline comet assay. Mutat Res. 2010; 683: 35–42.
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 17](#)
- 75
Garaj-Vrhovac V, Gajski G, Pažanin S, et al. Assessment of cytogenetic damage and oxidative stress in personnel occupationally exposed to the pulsed microwave radiation of marine radar equipment. Int J Hyg Environ Health. 2011; 214: 59–65.
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 4](#)
- 76
Hong R, Zhang Y, Liu Y, et al. Effects of extremely low frequency electromagnetic fields on DNA of testicular cells and sperm chromatin structure in mice. Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi. 2005; 23: 414–7. [Article in Chinese].
 - [PubMed](#)
- 77
Ivancsits S, Diem E, Pilger A, et al. Induction of DNA strand breaks by intermittent exposure to extremely-low-frequency electromagnetic fields in human diploid fibroblasts. Mutat Res. 2002; 519: 1–13.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 93](#)
- 78
Ivancsits S, Diem E, Jahn O, et al. Intermittent extremely low frequency electromagnetic fields cause DNA damage in a dose-dependent way. Int Arch Occup Environ Health. 2003; 76: 431–6.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 56](#)
- 79
Ivancsits S, Pilger A, Diem E, et al. Cell type-specific genotoxic effects of intermittent extremely low-frequency electromagnetic fields. Mutat Res. 2005; 583: 184–8.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),

- [Web of Science® Times Cited: 43](#)
- 80
Kesari KK, Behari J, Kumar S. Mutagenic response of 2.45 GHz radiation exposure on rat brain. Int J Radiat Biol. 2010; 86: 334–43.
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 17](#)
- 81
Lai H, Singh NP. Melatonin and a spin-trap compound block radiofrequency electromagnetic radiation-induced DNA strand breaks in rat brain cells. Bioelectromagnetics. 1997; 18: 446–54.
 Direct Link:
 - [Abstract](#)
 - [PDF\(203K\)](#)
 - [References](#)
 - [Web of Science® Times Cited: 108](#)
- 82
Lai H, Singh NP. Magnetic-field-induced DNA strand breaks in brain cells of the rat. Environ Health Perspect. 2004; 112: 687–94.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 96](#)
- 83
Lee JW, Kim MS, Kim YJ, et al. Genotoxic effects of 3 T magnetic resonance imaging in cultured human lymphocytes. Bioelectromagnetics. 2011; 32: 535–42.
 Direct Link:
 - [Abstract](#)
 - [Full Article \(HTML\)](#)
 - [PDF\(3714K\)](#)
 - [References](#)
 - [Web of Science® Times Cited: 3](#)
- 84
Paulraj R, Behari J. Single strand DNA breaks in rat brain cells exposed to microwave radiation. Mutat Res. 2006; 596: 76–80.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 43](#)
- 85
Romeo S, Zeni L, Sarti M, et al. DNA electrophoretic migration patterns change after exposure of Jurkat cells to a single intense nanosecond electric pulse. PLoS ONE. 2011; 6: e28419.
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 3](#),
 - [ADS](#)
- 86
Schwarz C, Kratochvil E, Pilger A, et al. Radiofrequency electromagnetic fields (UMTS, 1,950 MHz) induce genotoxic effects in vitro in human fibroblasts but not in lymphocytes. Int Arch Occup Environ Health. 2008; 81: 755–67.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 34](#)
- 87

- Svedenstål BM, Johanson KJ, Mattsson MO, et al. DNA damage, cell kinetics and ODC activities studied in CBA mice exposed to electromagnetic fields generated by transmission lines. *In Vivo*. 1999; 13: 507–13.
- 88
Svedenstål BM, Johanson KJ, Mild KH. DNA damage induced in brain cells of CBA mice exposed to magnetic fields. *In Vivo*. 1999; 13: 551–2.
 - 89
Trosić I, Pavčić I, Milković-Kraus S, et al. Effect of electromagnetic radiofrequency radiation on the rats' brain, liver and kidney cells measured by comet assay. *Coll Antropol*. 2011; 35: 1259–64.
 - 90
Burdak-Rothkamm S, Rothkamm K, Folkard M, et al. DNA and chromosomal damage in response to intermittent extremely low-frequency magnetic fields. *Mutat Res*. 2009; 672: 82–9.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 3](#)
 - 91
Fairbairn DW, O'Neill KL. The effect of electromagnetic field exposure on the formation of DNA single strand breaks in human cells. *Cell Mol Biol (Noisy-le-grand)*. 1994; 40: 561–7.
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 44](#)
 - 92
Fiorani M, Cantoni O, Sestili P, et al. Electric and/or magnetic field effects on DNA structure and function in cultured human cells. *Mutat Res*. 1992; 282: 25–9.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 43](#)
 - 93
Malyapa RS, Ahern EW, Straube WL, et al. Measurement of DNA damage after exposure to 2450 MHz electromagnetic radiation. *Radiat Res*. 1997; 148: 608–17.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 66](#)
 - 94
McNamee JP, Bellier PV, Chauhan V, et al. Evaluating DNA damage in rodent brain after acute 60 Hz magnetic-field exposure. *Radiat Res*. 2005; 164: 791–7.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 11](#)
 - 95
Scarfi MR, Sannino A, Perrotta A, et al. Evaluation of genotoxic effects in human fibroblasts after intermittent exposure to 50 Hz electromagnetic fields: a confirmatory study. *Radiat Res*. 2005; 164: 270–6.
 - [CrossRef](#),
 - [PubMed](#),
 - [Web of Science® Times Cited: 21](#)
 - 96
Stronati L, Testa A, Villani P, et al. Absence of genotoxicity in human blood cells exposed to 50 Hz magnetic fields as assessed by comet assay, chromosome aberration, micronucleus, and sister chromatid exchange analyses. *Bioelectromagnetics*. 2004; 25: 41–8.

Direct Link:

- [Abstract](#)
- [PDF\(113K\)](#)
- [References](#)
- [Web of Science® Times Cited: 25](#)

• 97

Testa A, Cordelli E, Stronati L, et al. Evaluation of genotoxic effect of low level 50 Hz magnetic fields on human blood cells using different cytogenetic assays. Bioelectromagnetics. 2004; 25: 613–9.

Direct Link:

- [Abstract](#)
- [PDF\(105K\)](#)
- [References](#)
- [Web of Science® Times Cited: 11](#)

• 98

Szabó G, Bährle S. Role of nitrosative stress and poly(ADP-ribose) polymerase activation in myocardial reperfusion injury. Curr Vasc Pharmacol. 2005; 3: 215–20.

- [CrossRef](#),
- [PubMed](#)

• 99

Moon HK, Yang ES, Park JW. Protection of peroxynitrite-induced DNA damage by dietary antioxidants. Arch Pharm Res. 2006; 29: 213–7.

- [CrossRef](#),
- [PubMed](#),
- [CAS](#),
- [Web of Science® Times Cited: 13](#)

• 100

Sakihama Y, Maeda M, Hashimoto M, et al. Beetroot betalain inhibits peroxynitrite-mediated tyrosine nitration and DNA strand damage. Free Radic Res. 2012; 46: 93–9.

- [CrossRef](#),
- [CAS](#),
- [Web of Science®](#)

• 101

Hybertson BM, Gao B, Bose SK, et al. Oxidative stress in health and disease: the therapeutic potential of Nrf2 activation. Mol Aspects Med. 2011; 32: 234–46.

- [CrossRef](#),
- [CAS](#),
- [Web of Science® Times Cited: 19](#)

• 102

Ridley AJ, Paterson HF, Johnston CL, et al. The small GTP-binding protein rac regulates growth factor-induced membrane ruffling. Cell. 1992; 70: 401–10.

- [CrossRef](#),
- [PubMed](#),
- [CAS](#),
- [Web of Science® Times Cited: 2665](#)

• 103

Wennström S, Hawkins P, Cooke F, et al. Activation of phosphoinositide 3-kinase is required for PDGF-stimulated membrane ruffling. Curr Biol. 1994; 4: 385–93.

- [CrossRef](#),
- [PubMed](#),
- [CAS](#),
- [Web of Science® Times Cited: 379](#)

• 104

Joucla S, Yvert B. Modeling of extracellular neural stimulation: from basic understanding to MEA-based applications. J Physiol Paris. 2012; 106: 146–58.

- [CrossRef](#),
 - [Web of Science® Times Cited: 2](#)
- 105
Pashut T, Wolfus S, Friedman A, et al. Mechanisms of magnetic stimulation of central nervous system neurons. PLoS Comput Biol. 2011; 7: e1002022.
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 7](#)
- 106
Fatemi-Ardekani A. Transcranial magnetic stimulation: physics, electrophysiology, and applications. Crit Rev Biomed Eng. 2008; 36: 375–412.
 - [CrossRef](#)
- 107
Silva S, Basser PJ, Miranda PC. Elucidating the mechanisms and loci of neuronal excitation by transcranial magnetic stimulation using a finite element model of a cortical sulcus. Clin Neurophysiol. 2008; 119: 2405–13.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 18](#)
- 108
Radman T, Ramos RL, Brumberg JC, et al. Role of cortical cell type and morphology in subthreshold and suprathreshold uniform electric field stimulation in vitro. Brain Stimul. 2009; 2: 215–28.
 - [CrossRef](#),
 - [Web of Science® Times Cited: 37](#)
- 109
Minelli TA, Balduzzo M, Milone FF, et al. Modeling cell dynamics under mobile phone radiation. Nonlinear cell dynamics under mobile phone radiation. Nonlinear Dynamics Psychol Life Sci. 2007; 11: 197–218.
 - [PubMed](#)
- 110
Saunders RD, Jefferys JGR. A neurobiological basis for ELF guidelines. Health Phys. 2007; 92: 596–603.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 15](#)
- 111
Havas M. Dirty electricity elevates blood sugar among electrically sensitive diabetics and may explain brittle diabetes. Electromagn Biol Med. 2008; 27: 135–46.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 5](#)
- 112
Havas M. Electromagnetic hypersensitivity: biological effects of dirty electricity with emphasis on diabetes and multiple sclerosis. Electromagn Biol Med. 2006; 25: 259–68.
 - [CrossRef](#),
 - [PubMed](#),
 - [Web of Science® Times Cited: 16](#)
- 113
de Vochta F. “Dirty electricity”: what, where, and should we care? J Expo Sci Environ Epidemiol. 2010; 20: 399–405.
 - [CrossRef](#),

- [Web of Science® Times Cited: 1](#)
- 114
Friedman J, Kraus S, Hauptman Y, et al. Mechanism of short-term ERK activation by electromagnetic fields at mobile phone frequencies. Biochem J. 2007; 405: 559–68.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 58](#)
- 115
Desai NR, Kesari KK, Agarwal A. Pathophysiology of cell phone radiation: oxidative stress and carcinogenesis with focus on the male reproductive system. Reproduct Biol Endocrinol. 2009; 7: 114. doi:[10.1186/1477-7827-7-114](https://doi.org/10.1186/1477-7827-7-114).
 - [CrossRef](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 15](#)
- 116
Wyatt CN, Weir EK, Peers C. Diphenylamine iodonium blocks K⁺ and Ca²⁺ currents in type I cells isolated from the rat carotid body. Neurosci Lett. 1994; 172: 63–6.
 - [CrossRef](#),
 - [PubMed](#),
 - [CAS](#),
 - [Web of Science® Times Cited: 28](#)